

Natural Deduction 1

Stéphane Devismes Pascal Lafourcade Michel Lévy
Jean-François Monin (jean-francois.monin@imag.fr)

Université Joseph Fourier, Grenoble I

February 6, 2015

Last course

- ▶ Davis-Putnam
- ▶ Complete Strategy

Homework : solution using Davis-Putnam

- ▶ (H1) : $p \Rightarrow \neg j \equiv \neg p \vee \neg j$
- ▶ (H2) : $\neg p \Rightarrow j \equiv p \vee j$
- ▶ (H3) : $j \Rightarrow m \equiv \neg j \vee m$
- ▶ (\neg C) : $\neg m \wedge \neg p$

Clauses : $\{\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p\}$

Homework : solution using Davis-Putnam

- ▶ (H1) : $p \Rightarrow \neg j \equiv \neg p \vee \neg j$
- ▶ (H2) : $\neg p \Rightarrow j \equiv p \vee j$
- ▶ (H3) : $j \Rightarrow m \equiv \neg j \vee m$
- ▶ (\neg C) : $\neg m \wedge \neg p$

Clauses : $\{\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p\}$

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

Homework : solution using Davis-Putnam

- ▶ (H1) : $p \Rightarrow \neg j \equiv \neg p \vee \neg j$
- ▶ (H2) : $\neg p \Rightarrow j \equiv p \vee j$
- ▶ (H3) : $j \Rightarrow m \equiv \neg j \vee m$
- ▶ (\neg C) : $\neg m \wedge \neg p$

Clauses : $\{\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p\}$

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

RE

Homework : solution using Davis-Putnam

- ▶ (H1) : $p \Rightarrow \neg j \equiv \neg p \vee \neg j$
- ▶ (H2) : $\neg p \Rightarrow j \equiv p \vee j$
- ▶ (H3) : $j \Rightarrow m \equiv \neg j \vee m$
- ▶ (\neg C) : $\neg m \wedge \neg p$

Clauses : $\{\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p\}$

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

RE

$$p \vee j, \neg j \vee m, \neg m, \neg p$$

Homework : solution using Davis-Putnam

- ▶ (H1) : $p \Rightarrow \neg j \equiv \neg p \vee \neg j$
- ▶ (H2) : $\neg p \Rightarrow j \equiv p \vee j$
- ▶ (H3) : $j \Rightarrow m \equiv \neg j \vee m$
- ▶ (\neg C) : $\neg m \wedge \neg p$

Clauses : $\{\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p\}$

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

RE

$$p \vee j, \neg j \vee m, \neg m, \neg p$$

UR on p

Homework : solution using Davis-Putnam

- ▶ (H1) : $p \Rightarrow \neg j \equiv \neg p \vee \neg j$
- ▶ (H2) : $\neg p \Rightarrow j \equiv p \vee j$
- ▶ (H3) : $j \Rightarrow m \equiv \neg j \vee m$
- ▶ (\neg C) : $\neg m \wedge \neg p$

Clauses : $\{\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p\}$

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

RE

$$p \vee j, \neg j \vee m, \neg m, \neg p$$

UR on p

$$j, \neg j \vee m, \neg m$$

Homework : solution using Davis-Putnam

- ▶ (H1) : $p \Rightarrow \neg j \equiv \neg p \vee \neg j$
- ▶ (H2) : $\neg p \Rightarrow j \equiv p \vee j$
- ▶ (H3) : $j \Rightarrow m \equiv \neg j \vee m$
- ▶ ($\neg C$) : $\neg m \wedge \neg p$

Clauses : $\{\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p\}$

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

RE

$$p \vee j, \neg j \vee m, \neg m, \neg p$$

UR on p

$$j, \neg j \vee m, \neg m$$

UR on m

Homework : solution using Davis-Putnam

- ▶ (H1) : $p \Rightarrow \neg j \equiv \neg p \vee \neg j$
- ▶ (H2) : $\neg p \Rightarrow j \equiv p \vee j$
- ▶ (H3) : $j \Rightarrow m \equiv \neg j \vee m$
- ▶ (\neg C) : $\neg m \wedge \neg p$

Clauses : $\{\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p\}$

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

RE

$$p \vee j, \neg j \vee m, \neg m, \neg p$$

UR on p

$$j, \neg j \vee m, \neg m$$

UR on m

$$j, \neg j$$

Homework : solution using Davis-Putnam

- ▶ (H1) : $p \Rightarrow \neg j \equiv \neg p \vee \neg j$
- ▶ (H2) : $\neg p \Rightarrow j \equiv p \vee j$
- ▶ (H3) : $j \Rightarrow m \equiv \neg j \vee m$
- ▶ ($\neg C$) : $\neg m \wedge \neg p$

Clauses : $\{\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p\}$

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

RE

$$p \vee j, \neg j \vee m, \neg m, \neg p$$

UR on p

$$j, \neg j \vee m, \neg m$$

UR on m

$$j, \neg j$$

UR on j

Homework : solution using Davis-Putnam

- ▶ (H1) : $p \Rightarrow \neg j \equiv \neg p \vee \neg j$
- ▶ (H2) : $\neg p \Rightarrow j \equiv p \vee j$
- ▶ (H3) : $j \Rightarrow m \equiv \neg j \vee m$
- ▶ ($\neg C$) : $\neg m \wedge \neg p$

Clauses : $\{\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p\}$

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

RE

$$p \vee j, \neg j \vee m, \neg m, \neg p$$

UR on p

$$j, \neg j \vee m, \neg m$$

UR on m

$$j, \neg j$$

UR on j

⊥

Homework : solution using complete strategy

 Δ_{i+1}

- ▶ Construct all the resolvents of Δ_i and of $\Delta_i \cup \Theta_i$
- ▶ Reduce this set
- ▶ Remove the new resolvents including a clause of $\Delta_i \cup \Theta_i$

Θ_{i+1} : Remove from $\Delta_i \cup \Theta_i$ the clauses including a clause of Δ_{i+1} .

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

Homework : solution using complete strategy

 Δ_{i+1}

- ▶ Construct all the resolvents of Δ_i and of $\Delta_i \cup \Theta_i$
- ▶ Reduce this set
- ▶ Remove the new resolvents including a clause of $\Delta_i \cup \Theta_i$

 Θ_{i+1} : Remove from $\Delta_i \cup \Theta_i$ the clauses including a clause of Δ_{i+1} .

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

k	Δ_k	Θ_k	$\Delta_k \cup \Theta_k$	Resol. of Δ_k and $\Delta_k \cup \Theta_k$
0	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	\emptyset	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	
1				
2				
3				

Homework : solution using complete strategy

 Δ_{i+1}

- ▶ Construct all the resolvents of Δ_i and of $\Delta_i \cup \Theta_i$
- ▶ Reduce this set
- ▶ Remove the new resolvents including a clause of $\Delta_i \cup \Theta_i$

 Θ_{i+1} : Remove from $\Delta_i \cup \Theta_i$ the clauses including a clause of Δ_{i+1} .

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

k	Δ_k	Θ_k	$\Delta_k \cup \Theta_k$	Resol. of Δ_k and $\Delta_k \cup \Theta_k$
0	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	\emptyset	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	j (1,4) $\neg j$ (2,3) $p \vee m$ (1,2)
1				
2				
3				

Homework : solution using complete strategy

 Δ_{i+1}

- ▶ Construct all the resolvents of Δ_i and of $\Delta_i \cup \Theta_i$
- ▶ Reduce this set
- ▶ Remove the new resolvents including a clause of $\Delta_i \cup \Theta_i$

 Θ_{i+1} : Remove from $\Delta_i \cup \Theta_i$ the clauses including a clause of Δ_{i+1} .

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

k	Δ_k	Θ_k	$\Delta_k \cup \Theta_k$	Resol. of Δ_k and $\Delta_k \cup \Theta_k$
0	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	\emptyset	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	j (1,4) $\neg j$ (2,3) $p \vee m$ (1,2)
1	5) j 6) $\neg j$ 7) $p \vee m$			
2				
3				

Homework : solution using complete strategy

 Δ_{i+1}

- ▶ Construct all the resolvents of Δ_i and of $\Delta_i \cup \Theta_i$
- ▶ Reduce this set
- ▶ Remove the new resolvents including a clause of $\Delta_i \cup \Theta_i$

 Θ_{i+1} : Remove from $\Delta_i \cup \Theta_i$ the clauses including a clause of Δ_{i+1} .

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

k	Δ_k	Θ_k	$\Delta_k \cup \Theta_k$	Resol. of Δ_k and $\Delta_k \cup \Theta_k$
0	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	\emptyset	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	j (1,4) $\neg j$ (2,3) $p \vee m$ (1,2)
1	5) j 6) $\neg j$ 7) $p \vee m$	$\neg m, \neg p$		
2				
3				

Homework : solution using complete strategy

 Δ_{i+1}

- ▶ Construct all the resolvents of Δ_i and of $\Delta_i \cup \Theta_i$
- ▶ Reduce this set
- ▶ Remove the new resolvents including a clause of $\Delta_i \cup \Theta_i$

 Θ_{i+1} : Remove from $\Delta_i \cup \Theta_i$ the clauses including a clause of Δ_{i+1} .

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

k	Δ_k	Θ_k	$\Delta_k \cup \Theta_k$	Resol. of Δ_k and $\Delta_k \cup \Theta_k$
0	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	\emptyset	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	j (1,4) $\neg j$ (2,3) $p \vee m$ (1,2)
1	5) j 6) $\neg j$ 7) $p \vee m$	$\neg m, \neg p$	5) j , 6) $\neg j$, 7) $p \vee m$ 3) $\neg m$, 4) $\neg p$	
2				
3				

Homework : solution using complete strategy

 Δ_{i+1}

- ▶ Construct all the resolvents of Δ_i and of $\Delta_i \cup \Theta_i$
- ▶ Reduce this set
- ▶ Remove the new resolvents including a clause of $\Delta_i \cup \Theta_i$

 Θ_{i+1} : Remove from $\Delta_i \cup \Theta_i$ the clauses including a clause of Δ_{i+1} .

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

k	Δ_k	Θ_k	$\Delta_k \cup \Theta_k$	Resol. of Δ_k and $\Delta_k \cup \Theta_k$
0	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	\emptyset	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	j (1,4) $\neg j$ (2,3) $p \vee m$ (1,2)
1	5) j 6) $\neg j$ 7) $p \vee m$	$\neg m, \neg p$	5) j , 6) $\neg j$, 7) $p \vee m$ 3) $\neg m$, 4) $\neg p$	m (4,7) p (7,3) \perp (5,6)
2				
3				

Homework : solution using complete strategy

 Δ_{i+1}

- ▶ Construct all the resolvents of Δ_i and of $\Delta_i \cup \Theta_i$
- ▶ Reduce this set
- ▶ Remove the new resolvents including a clause of $\Delta_i \cup \Theta_i$

 Θ_{i+1} : Remove from $\Delta_i \cup \Theta_i$ the clauses including a clause of Δ_{i+1} .

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

k	Δ_k	Θ_k	$\Delta_k \cup \Theta_k$	Resol. of Δ_k and $\Delta_k \cup \Theta_k$
0	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	\emptyset	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	j (1,4) $\neg j$ (2,3) $p \vee m$ (1,2)
1	5) j 6) $\neg j$ 7) $p \vee m$	$\neg m, \neg p$	5) j , 6) $\neg j$, 7) $p \vee m$ 3) $\neg m$, 4) $\neg p$	m (4,7) p (7,3) \perp (5,6)
2	\perp			
3				

Homework : solution using complete strategy

 Δ_{i+1}

- ▶ Construct all the resolvents of Δ_i and of $\Delta_i \cup \Theta_i$
- ▶ Reduce this set
- ▶ Remove the new resolvents including a clause of $\Delta_i \cup \Theta_i$

 Θ_{i+1} : Remove from $\Delta_i \cup \Theta_i$ the clauses including a clause of Δ_{i+1} .

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

k	Δ_k	Θ_k	$\Delta_k \cup \Theta_k$	Resol. of Δ_k and $\Delta_k \cup \Theta_k$
0	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	\emptyset	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	j (1,4) $\neg j$ (2,3) $p \vee m$ (1,2)
1	5) j 6) $\neg j$ 7) $p \vee m$	$\neg m, \neg p$	5) j , 6) $\neg j$, 7) $p \vee m$ 3) $\neg m$, 4) $\neg p$	m (4,7) p (7,3) \perp (5,6)
2	\perp	\emptyset		
3				

Homework : solution using complete strategy

 Δ_{i+1}

- ▶ Construct all the resolvents of Δ_i and of $\Delta_i \cup \Theta_i$
- ▶ Reduce this set
- ▶ Remove the new resolvents including a clause of $\Delta_i \cup \Theta_i$

 Θ_{i+1} : Remove from $\Delta_i \cup \Theta_i$ the clauses including a clause of Δ_{i+1} .

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

k	Δ_k	Θ_k	$\Delta_k \cup \Theta_k$	Resol. of Δ_k and $\Delta_k \cup \Theta_k$
0	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	\emptyset	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	j (1,4) $\neg j$ (2,3) $p \vee m$ (1,2)
1	5) j 6) $\neg j$ 7) $p \vee m$	$\neg m, \neg p$	5) j , 6) $\neg j$, 7) $p \vee m$ 3) $\neg m$, 4) $\neg p$	m (4,7) p (7,3) \perp (5,6)
2	\perp	\emptyset	\perp	
3				

Homework : solution using complete strategy

 Δ_{i+1}

- ▶ Construct all the resolvents of Δ_i and of $\Delta_i \cup \Theta_i$
- ▶ Reduce this set
- ▶ Remove the new resolvents including a clause of $\Delta_i \cup \Theta_i$

 Θ_{i+1} : Remove from $\Delta_i \cup \Theta_i$ the clauses including a clause of Δ_{i+1} .

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

k	Δ_k	Θ_k	$\Delta_k \cup \Theta_k$	Resol. of Δ_k and $\Delta_k \cup \Theta_k$
0	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	\emptyset	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	j (1,4) $\neg j$ (2,3) $p \vee m$ (1,2)
1	5) j 6) $\neg j$ 7) $p \vee m$	$\neg m, \neg p$	5) j , 6) $\neg j$, 7) $p \vee m$ 3) $\neg m$, 4) $\neg p$	m (4,7) p (7,3) \perp (5,6)
2	\perp	\emptyset	\perp	\emptyset
3				

Homework : solution using complete strategy

 Δ_{i+1}

- ▶ Construct all the resolvents of Δ_i and of $\Delta_i \cup \Theta_i$
- ▶ Reduce this set
- ▶ Remove the new resolvents including a clause of $\Delta_i \cup \Theta_i$

 Θ_{i+1} : Remove from $\Delta_i \cup \Theta_i$ the clauses including a clause of Δ_{i+1} .

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

k	Δ_k	Θ_k	$\Delta_k \cup \Theta_k$	Resol. of Δ_k and $\Delta_k \cup \Theta_k$
0	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	\emptyset	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	j (1,4) $\neg j$ (2,3) $p \vee m$ (1,2)
1	5) j 6) $\neg j$ 7) $p \vee m$	$\neg m, \neg p$	5) j , 6) $\neg j$, 7) $p \vee m$ 3) $\neg m$, 4) $\neg p$	m (4,7) p (7,3) \perp (5,6)
2	\perp	\emptyset	\perp	\emptyset
3	\emptyset			

Homework : solution using complete strategy

 Δ_{i+1}

- ▶ Construct all the resolvents of Δ_i and of $\Delta_i \cup \Theta_i$
- ▶ Reduce this set
- ▶ Remove the new resolvents including a clause of $\Delta_i \cup \Theta_i$

 Θ_{i+1} : Remove from $\Delta_i \cup \Theta_i$ the clauses including a clause of Δ_{i+1} .

$$\neg p \vee \neg j, p \vee j, \neg j \vee m, \neg m, \neg p$$

k	Δ_k	Θ_k	$\Delta_k \cup \Theta_k$	Resol. of Δ_k and $\Delta_k \cup \Theta_k$
0	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	\emptyset	1) $p \vee j$ 2) $\neg j \vee m$ 3) $\neg m$ 4) $\neg p$	j (1,4) $\neg j$ (2,3) $p \vee m$ (1,2)
1	5) j 6) $\neg j$ 7) $p \vee m$	$\neg m, \neg p$	5) j , 6) $\neg j$, 7) $p \vee m$ 3) $\neg m$, 4) $\neg p$	m (4,7) p (7,3) \perp (5,6)
2	\perp	\emptyset	\perp	\emptyset
3	\emptyset	\perp		

Plan

Introduction

Rules

Natural deduction proofs

Correctness

Completeness

Tactics

Conclusion

Plan

Introduction

Rules

Natural deduction proofs

Correctness

Completeness

Tactics

Conclusion

Intuition

When we write proofs in math courses,
when we decompose a reasoning in elementary obvious steps,
we practice **natural deduction**.

The natural deduction ND

History : In 1934 **Gerhard Gentzen**

Introduced two models of ND for classical logic :

- ▶ **NK** : a proof is a tree of formulas.
- ▶ **LK** : a proof is a tree of sequents.

NJ et **LJ** for intuitionistic logic.



Here, **yet another** presentation of natural deduction.

Resolution vs. Natural deduction

A proof by **resolution** is a **list of clauses**.

In **natural deduction**, during a proof,
we can add and remove hypotheses,

Resolution vs. Natural deduction

A proof by **resolution** is a **list of clauses**.

In **natural deduction**, during a proof,
we can add and remove hypotheses,

Hence a more complex definition of a proof.

Abbreviations

Negation and equivalence are **abbreviations** defined as :

- ▶ \top abbreviates to $\perp \Rightarrow \perp$.
- ▶ $\neg A$ abbreviates to $A \Rightarrow \perp$.
- ▶ $A \Leftrightarrow B$ abbreviates to $(A \Rightarrow B) \wedge (B \Rightarrow A)$.

Abbreviations

Negation and equivalence are **abbreviations** defined as :

- ▶ \top abbreviates to $\perp \Rightarrow \perp$.
- ▶ $\neg A$ abbreviates to $A \Rightarrow \perp$.
- ▶ $A \Leftrightarrow B$ abbreviates to $(A \Rightarrow B) \wedge (B \Rightarrow A)$.

Two formulae are considered to be **equal**, if the formulas obtained by removing the abbreviations are identical.

Abbreviations

Negation and equivalence are **abbreviations** defined as :

- ▶ \top abbreviates to $\perp \Rightarrow \perp$.
- ▶ $\neg A$ abbreviates to $A \Rightarrow \perp$.
- ▶ $A \Leftrightarrow B$ abbreviates to $(A \Rightarrow B) \wedge (B \Rightarrow A)$.

Two formulae are considered to be **equal**, if the formulas obtained by removing the abbreviations are identical.

For example, the formulae $\neg\neg a$, $\neg a \Rightarrow \perp$ and $(a \Rightarrow \perp) \Rightarrow \perp$ are equal.

Abbreviations

Negation and equivalence are **abbreviations** defined as :

- ▶ \top abbreviates to $\perp \Rightarrow \perp$.
- ▶ $\neg A$ abbreviates to $A \Rightarrow \perp$.
- ▶ $A \Leftrightarrow B$ abbreviates to $(A \Rightarrow B) \wedge (B \Rightarrow A)$.

Two formulae are considered to be **equal**, if the formulas obtained by removing the abbreviations are identical.

For example, the formulae $\neg\neg a$, $\neg a \Rightarrow \perp$ and $(a \Rightarrow \perp) \Rightarrow \perp$ are equal.

Two equal formulae are equivalent !

Plan

Introduction

Rules

Natural deduction proofs

Correctness

Completeness

Tactics

Conclusion

Rule

Definition 3.1.1

A **rule** consists of formulas called **premises** (sometimes called hypotheses) H_1, \dots, H_n and of a unique **conclusion**.

Premises are written above a line and the conclusion below this line.

The name of the rule is written at the same level as the line.

$$\frac{H_1 \dots H_n}{C} R$$

Fundamental rule of Natural Deduction

Fundamental rule of Natural Deduction

Implies-introduction :

In order to prove $A \Rightarrow B$,
just derive B with the additional hypothesis A .

Classification of rules

Classification of rules

- ▶ **Introduction rules** for introducing a connective in the conclusion.

Classification of rules

- ▶ **Introduction rules** for introducing a connective in the conclusion.
- ▶ **Elimination rules** for removing a connective from the premises.

Classification of rules

- ▶ **Introduction rules** for introducing a connective in the conclusion.
- ▶ **Elimination rules** for removing a connective from the premises.
- ▶ + **two special rules**

Rules (system NK of Gentzen)

Table 3.1

Introduction	Élimination
$\begin{array}{c} [A] \\ \vdots \\ B \\ \hline A \Rightarrow B \end{array} \Rightarrow I$	$\frac{A \quad A \Rightarrow B}{B} \Rightarrow E$
$\frac{A \quad B}{A \wedge B} \wedge I$	$\frac{A \wedge B}{A} \wedge E1$
	$\frac{A \wedge B}{B} \wedge E2$
$\frac{A}{A \vee B} \vee I1$	$\frac{A \vee B \quad A \Rightarrow C \quad B \Rightarrow C}{C} \vee E$
$\frac{A}{B \vee A} \vee I2$	
Rule about false	
$\frac{}{A} \perp \text{ Eq}$	
Reductio ad absurdum	
$\frac{\neg \neg A}{A} \text{ RAA}$	

[A] means that A is a hypothesis

Plan

Introduction

Rules

Natural deduction proofs

Correctness

Completeness

Tactics

Conclusion

Principle

- ▶ A proof is a sequence of lines ; lines are derived from previous lines using the rules.

Principle

- ▶ A proof is a sequence of lines ; lines are derived from previous lines using the rules.
- ▶ An additional hypothesis A is assumed by the line : Suppose A .

Principle

- ▶ A proof is a sequence of lines ; lines are derived from previous lines using the rules.
- ▶ An additional hypothesis A is assumed by the line : Suppose A .
- ▶ The last hypothesis is removed A by the line : Hence $A \Rightarrow B$.

Principle

- ▶ A proof is a sequence of lines ; lines are derived from previous lines using the rules.
- ▶ An additional hypothesis A is assumed by the line : Suppose A .
- ▶ The last hypothesis is removed A by the line : Hence $A \Rightarrow B$.
- ▶ This line is **the rule of implies-introduction**.

$$\frac{\begin{array}{c} [A] \\ \vdots \\ B \end{array}}{A \Rightarrow B} \Rightarrow I$$

Proof line

Definition 3.1.2

A proof **line** is of one of the three following forms :

- ▶ Suppose **formula**,
- ▶ **formula**,
- ▶ Hence **formula**.

Proof line

Definition 3.1.2

A proof **line** is of one of the three following forms :

- ▶ Suppose **formula**,
- ▶ **formula**,
- ▶ Hence **formula**.

Examples :

- ▶ Suppose $A \wedge B$
- ▶ A
- ▶ Hence $A \wedge B \Rightarrow A$

Scratch proof

Definition 3.1.3

A **scratch** proof is a sequence of lines such that, in every prefix of the sequence, the number of lines starting with the word `Suppose` is at least equal to the number of lines starting with the word `Hence`.

Scratch proof

Definition 3.1.3

A **scratch** proof is a sequence of lines such that, in every prefix of the sequence, the number of lines starting with the word *Suppose* is at least equal to the number of lines starting with the word *Hence*.

Example 3.1.4

number	line
1	Suppose a
2	$a \vee b$
3	Hence $a \Rightarrow a \vee b$
4	Hence $\neg a$
5	Suppose b

Scratch proof : examples

Where are the scratches ?

num	line
1	Suppose $a \wedge b$
2	b
3	$b \vee c$
4	Hence $a \wedge b \Rightarrow b \vee c$
5	Hence $\neg a$
6	Suppose b

num	line
1	Suppose a
2	$a \vee b$
3	Hence $a \Rightarrow a \vee b$
4	Suppose b

num	line
1	Suppose a
2	$a \vee b$
3	Hence $a \Rightarrow a \vee b$
4	Suppose b
5	Hence $\neg a$

Context (1/2)

- ▶ Each line of a scratch proof contains a **context**
- ▶ A **context** is the sequence of hypotheses previously introduced in lines `Suppose` and not removed in lines `Hence`.

Context (1/2)

- ▶ Each line of a scratch proof contains a **context**
- ▶ A **context** is the sequence of hypotheses previously introduced in lines *Suppose* and not removed in lines *Hence*.

Example 3.1.6 :

context	number	line	rule
1	1	Suppose a	
1,2	2	Suppose b	
1,2	3	$a \wedge b$	$\wedge I$ 1,2
1	4	Hence $b \Rightarrow a \wedge b$	$\Rightarrow I$ 2,3
1,5	5	Suppose e	

Context (2/2)

Definition 3.1.5

Lines of a scratch proof are numbered from 1 to n . For i between 1 and n , the list of formulae Γ_i is the **context** of the line i . The list Γ_0 is empty and the lists of formulae Γ_i are defined as :

- ▶ If the line i is « Suppose A », then $\Gamma_i = \Gamma_{i-1}, i$.
- ▶ If the line i is « A » then $\Gamma_i = \Gamma_{i-1}$
- ▶ If the line i is « Hence A » then Γ_i is obtained by eliminated the last formula of Γ_{i-1}

The list Γ_i is the **context** of the line i .

The context of a formula represents the hypotheses from which it has been derived.

Example of context

Give the **context** of the following proof scratch :

context	number	line
	1	Suppose a
	2	$a \vee b$
	3	Hence $a \Rightarrow a \vee b$
	4	Suppose b
	5	Hence b

Example of context

Give the **context** of the following proof scratch :

context	number	line
1	1	Suppose a
1	2	$a \vee b$
	3	Hence $a \Rightarrow a \vee b$
4	4	Suppose b
	5	Hence b

Usable formulae (1/3)

Definition 3.1.7

- ▶ A formula appearing on a line of a scratch proof is the **conclusion** of the line.
- ▶ The conclusion of a line is **usable** as long as its context (i.e., the context from which it has been derived) is present.

Usable formulae (2/3)

Example 3.1.8

context	number	line
1	1	Suppose a
1	2	$a \vee b$
	3	Hence $a \Rightarrow b$
	4	a
	5	$b \vee a$

The conclusion of line 2 is usable over the line 2, and not beyond, as on line 3, the hypothesis from which it has been derived is eliminated.

Usable formulae (3/3)

Give the lines for which lines 1 and 3 are **usable** in the following example :

context	number	line
1	1	Suppose a
1,2	2	Suppose b
1,2	3	c
1	4	Hence d
1,5	5	Suppose e

Definition of a Proof

Definition 3.1.9

Let Γ a set of formulae, a **proof in the environment** Γ is a scratch proof having the following properties :

1. For every line $\ll \text{Hence } A \gg$, the formula A is $B \Rightarrow C$, where B is the last formula from the context of the previous line and where C is a usable formula for the previous line or is a member of the environment Γ .
2. For every line $\ll A \gg$, the formula A is the conclusion of a rule (other than the rule of implies-introduction) whose premises are usable on the previous line, or are members of the environment Γ .

Proof of formulae

Definition 3.1.10

A **proof of the formulae** A within the environment Γ is :

- ▶ either the empty proof when A is element of Γ ,
- ▶ or a proof whose last line is A with an empty context.

Proof of formulae

Definition 3.1.10

A **proof of the formulae** A within the environment Γ is :

- ▶ either the empty proof when A is element of Γ ,
- ▶ or a proof whose last line is A with an empty context.

We note :

- ▶ $\Gamma \vdash A$ the fact that there is a proof of A within the environment Γ ,
- ▶ $\Gamma \vdash P : A$ the fact that P is a proof of A within the environment Γ .
- ▶ When the environment is empty, we abbreviate $\emptyset \vdash A$ by $\vdash A$.
- ▶ When we ask for a proof of a formula without indicating the environment, we suppose that $\Gamma = \emptyset$.

First Example (exemple 3.1.11)

Let us prove $(a \Rightarrow b) \Rightarrow (\neg b \Rightarrow \neg a)$.

context	number	proof	justification
---------	--------	-------	---------------

First Example (exemple 3.1.11)

Let us prove $(a \Rightarrow b) \Rightarrow (\neg b \Rightarrow \neg a)$.

context	number	proof	justification
1	1	Suppose $a \Rightarrow b$	

First Example (exemple 3.1.11)

Let us prove $(a \Rightarrow b) \Rightarrow (\neg b \Rightarrow \neg a)$.

context	number	proof	justification
1	1	Suppose $a \Rightarrow b$	
1,2	2	Suppose $\neg b$	

First Example (exemple 3.1.11)

Let us prove $(a \Rightarrow b) \Rightarrow (\neg b \Rightarrow \neg a)$.

context	number	proof	justification
1	1	Suppose $a \Rightarrow b$	
1,2	2	Suppose $\neg b$	
1,2,3	3	Suppose a	

First Example (exemple 3.1.11)

Let us prove $(a \Rightarrow b) \Rightarrow (\neg b \Rightarrow \neg a)$.

context	number	proof	justification
1	1	Suppose $a \Rightarrow b$	
1,2	2	Suppose $\neg b$	
1,2,3	3	Suppose a	
1,2,3	4	b	$\Rightarrow E$ 1, 3

First Example (exemple 3.1.11)

Let us prove $(a \Rightarrow b) \Rightarrow (\neg b \Rightarrow \neg a)$.

context	number	proof	justification
1	1	Suppose $a \Rightarrow b$	
1,2	2	Suppose $\neg b$	
1,2,3	3	Suppose a	
1,2,3	4	b	$\Rightarrow E$ 1, 3
1,2,3	5	\perp	$\Rightarrow E$ 2, 4

First Example (exemple 3.1.11)

Let us prove $(a \Rightarrow b) \Rightarrow (\neg b \Rightarrow \neg a)$.

context	number	proof	justification
1	1	Suppose $a \Rightarrow b$	
1,2	2	Suppose $\neg b$	
1,2,3	3	Suppose a	
1,2,3	4	b	$\Rightarrow E$ 1, 3
1,2,3	5	\perp	$\Rightarrow E$ 2, 4
1,2	6	Hence $\neg a$	$\Rightarrow I$ 3, 5

First Example (exemple 3.1.11)

Let us prove $(a \Rightarrow b) \Rightarrow (\neg b \Rightarrow \neg a)$.

context	number	proof	justification
1	1	Suppose $a \Rightarrow b$	
1,2	2	Suppose $\neg b$	
1,2,3	3	Supposons a	
1,2,3	4	b	$\Rightarrow E$ 1, 3
1,2,3	5	\perp	$\Rightarrow E$ 2, 4
1,2	6	Hence $\neg a$	$\Rightarrow I$ 3, 5
1	7	Hence $\neg b \Rightarrow \neg a$	$\Rightarrow I$ 2, 6

First Example (exemple 3.1.11)

Let us prove $(a \Rightarrow b) \Rightarrow (\neg b \Rightarrow \neg a)$.

context	number	proof	justification
1	1	Suppose $a \Rightarrow b$	
1,2	2	Suppose $\neg b$	
1,2,3	3	Suppose a	
1,2,3	4	b	$\Rightarrow E$ 1, 3
1,2,3	5	\perp	$\Rightarrow E$ 2, 4
1,2	6	Hence $\neg a$	$\Rightarrow I$ 3, 5
1	7	Hence $\neg b \Rightarrow \neg a$	$\Rightarrow I$ 2, 6
	8	Hence $(a \Rightarrow b) \Rightarrow (\neg b \Rightarrow \neg a)$	$\Rightarrow I$ 1, 7

Proofs with abbreviations vs. without abbreviations

cont.	n.	proof with abbreviation	proof without abbreviation	just.
1	1	Suppose $a \Rightarrow b$	Suppose $a \Rightarrow b$	
1,2	2	Suppose $\neg b$	Suppose $b \Rightarrow \perp$	
1,2,3	3	Suppose a	Suppose a	
1,2,3	4	b	b	$\Rightarrow E 1, 3$
1,2,3	5	\perp	\perp	$\Rightarrow E 2, 4$
1,2	6	Hence $\neg a$	Hence $a \Rightarrow \perp$	$\Rightarrow I 3, 5$
1	7	Hence $\neg b \Rightarrow \neg a$	Hence $(b \Rightarrow \perp) \Rightarrow (a \Rightarrow \perp)$	$\Rightarrow I 2, 6$
	8	Hence $(a \Rightarrow b) \Rightarrow (\neg b \Rightarrow \neg a)$	Hence $(a \Rightarrow b) \Rightarrow ((b \Rightarrow \perp) \Rightarrow (a \Rightarrow \perp))$	$\Rightarrow I 1, 7$

Tree (example 3.1.11)

$$\begin{array}{c}
 \frac{\frac{\frac{(1)a \Rightarrow b \quad (3)\phi}{(4)b} \Rightarrow E}{(2)\neg b} \Rightarrow E}{(5)\perp} \Rightarrow E \\
 \frac{(5)\perp}{(6)\neg a} \Rightarrow I[3] \\
 \frac{(6)\neg a}{(7)\neg b \Rightarrow \neg a} \Rightarrow I[2] \\
 \frac{(7)\neg b \Rightarrow \neg a}{(8)(a \Rightarrow b) \Rightarrow (\neg b \Rightarrow \neg a)} \Rightarrow I[1]
 \end{array}$$

context	number	proof	justification
1	1	Suppose $a \Rightarrow b$	
1,2	2	Suppose $\neg b$	
1,2,3	3	Suppose a	
1,2,3	4	b	$\Rightarrow E$ 1, 3
1,2,3	5	\perp	$\Rightarrow E$ 2, 4
1,2	6	Hence $\neg a$	$\Rightarrow I$ 3, 5
1	7	Hence $\neg b \Rightarrow \neg a$	$\Rightarrow I$ 2, 6
	8	Hence $(a \Rightarrow b) \Rightarrow (\neg b \Rightarrow \neg a)$	$\Rightarrow I$ 1, 7

Second Example

Prove $a \wedge \neg a \Rightarrow b$.

context	number	proof	justification
---------	--------	-------	---------------

Second Example

Prove $a \wedge \neg a \Rightarrow b$.

context	number	proof	justification
1	1	Suppose $a \wedge \neg a$	

Second Example

Prove $a \wedge \neg a \Rightarrow b$.

context	number	proof	justification
1	1	Suppose $a \wedge \neg a$	
1	2	a	$\wedge E1$ 1

Second Example

Prove $a \wedge \neg a \Rightarrow b$.

context	number	proof	justification
1	1	Suppose $a \wedge \neg a$	
1	2	a	$\wedge E1$ 1
1	3	$\neg a$	$\wedge E2$ 1

Second Example

Prove $a \wedge \neg a \Rightarrow b$.

context	number	proof	justification
1	1	Suppose $a \wedge \neg a$	
1	2	a	$\wedge E1$ 1
1	3	$\neg a$	$\wedge E2$ 1
1	4	\perp	$\Rightarrow E$ 2,3

Second Example

Prove $a \wedge \neg a \Rightarrow b$.

context	number	proof	justification
1	1	Suppose $a \wedge \neg a$	
1	2	a	$\wedge E1$ 1
1	3	$\neg a$	$\wedge E2$ 1
1	4	\perp	$\Rightarrow E$ 2,3
1	5	b	Efq 4

Second Example

Prove $a \wedge \neg a \Rightarrow b$.

context	number	proof	justification
1	1	Suppose $a \wedge \neg a$	
1	2	a	$\wedge E1$ 1
1	3	$\neg a$	$\wedge E2$ 1
1	4	\perp	$\Rightarrow E$ 2,3
1	5	b	$Efq4$
	6	Hence $a \wedge \neg a \Rightarrow b$	$\Rightarrow I$ 1,5

Proofs with abbreviations vs. without abbreviation (2/2)

contexte	number	proof with abbreviation	proof without abbreviation	justification
1	1	Supposons $a \wedge \neg a$	Supposons $a \wedge (a \Rightarrow \perp)$	
1	2	a	a	$\wedge E1$ 1
1	3	$\neg a$	$a \Rightarrow \perp$	$\wedge E2$ 1
1	4	\perp	\perp	$\Rightarrow E$ 2,3
1	5	b	b	$Efq4$
	6	Donc $a \wedge \neg a \Rightarrow b$	Donc $a \wedge (a \Rightarrow \perp) \Rightarrow b$	$\Rightarrow I$ 1,5

Third Example

Prove $\neg A$ in the environment $\neg(A \vee B)$

environment			
reference		formula	
i		$\neg(A \vee B)$	
context	number	proof	justification

Third Example

Prove $\neg A$ in the environment $\neg(A \vee B)$

environment			
reference		formula	
i		$\neg(A \vee B)$	
context	number	proof	justification
1	1	Suppose A	

Third Example

Prove $\neg A$ in the environment $\neg(A \vee B)$

environment			
reference		formula	
i		$\neg(A \vee B)$	
context	number	proof	justification
1	1	Suppose A	
1	2	$A \vee B$	$\vee I$ 1

Third Example

Prove $\neg A$ in the environment $\neg(A \vee B)$

environment			
reference		formula	
i		$\neg(A \vee B)$	
context	number	proof	justification
1	1	Suppose A	
1	2	$A \vee B$	$\vee I$ 1
1	3	\perp	$\Rightarrow E$ $i, 2$

Third Example

Prove $\neg A$ in the environment $\neg(A \vee B)$

environment			
reference		formula	
i		$\neg(A \vee B)$	
context	number	proof	justification
1	1	Suppose A	
1	2	$A \vee B$	$\vee I$ 1
1	3	\perp	$\Rightarrow E$ $i, 2$
	4	Hence $\neg A$	$\Rightarrow I$ 1, 3

Forth exemple (example 3.1.12)

Prove $\neg A \vee B$ in the environment $A \Rightarrow B$.

environment			
reference		formule	
i		$A \Rightarrow B$	
context	number	proof	justification

Forth exemple (example 3.1.12)

Prove $\neg A \vee B$ in the environment $A \Rightarrow B$.

environment			
reference		formula	
i		$A \Rightarrow B$	
context	number	proof	justification
1	1	Suppose $\neg(\neg A \vee B)$	

Forth exemple (example 3.1.12)

Prove $\neg A \vee B$ in the environment $A \Rightarrow B$.

environment			
reference		formula	
i		$A \Rightarrow B$	
context	number	proof	justification
1	1	Suppose $\neg(\neg A \vee B)$	
1,2	2	Suppose A	

Forth exemple (example 3.1.12)

Prove $\neg A \vee B$ in the environment $A \Rightarrow B$.

environment			
reference		formula	
i		$A \Rightarrow B$	
context	number	proof	justification
1	1	Suppose $\neg(\neg A \vee B)$	
1,2	2	Suppose A	
1,2	3	B	$\Rightarrow E\ i, 2$

Forth exemple (example 3.1.12)

Prove $\neg A \vee B$ in the environment $A \Rightarrow B$.

environment			
reference		formula	
i		$A \Rightarrow B$	
context	number	proof	justification
1	1	Suppose $\neg(\neg A \vee B)$	
1,2	2	Suppose A	
1,2	3	B	$\Rightarrow E$ $i, 2$
1,2	4	$\neg A \vee B$	$\vee I$ $2, 3$

Forth exemple (example 3.1.12)

Prove $\neg A \vee B$ in the environment $A \Rightarrow B$.

environment			
reference		formula	
i		$A \Rightarrow B$	
context	number	proof	justification
1	1	Suppose $\neg(\neg A \vee B)$	
1,2	2	Suppose A	
1,2	3	B	$\Rightarrow E$ $i, 2$
1,2	4	$\neg A \vee B$	$\vee I$ $2, 3$
1,2	5	\perp	$\Rightarrow E$ $1, 4$

Forth exemple (example 3.1.12)

Prove $\neg A \vee B$ in the environment $A \Rightarrow B$.

environment			
reference		formula	
<i>i</i>		$A \Rightarrow B$	
context	number	proof	justification
1	1	Suppose $\neg(\neg A \vee B)$	
1,2	2	Suppose A	
1,2	3	B	$\Rightarrow E\ i, 2$
1,2	4	$\neg A \vee B$	$\vee I\ 2\ 3$
1,2	5	\perp	$\Rightarrow E\ 1, 4$
1	6	Hence $\neg A$	$\Rightarrow I\ 2, 5$

Forth exemple (example 3.1.12)

Prove $\neg A \vee B$ in the environment $A \Rightarrow B$.

environment			
reference		formula	
<i>i</i>		$A \Rightarrow B$	
context	number	proof	justification
1	1	Suppose $\neg(\neg A \vee B)$	
1,2	2	Suppose A	
1,2	3	B	$\Rightarrow E$ 1, 2
1,2	4	$\neg A \vee B$	$\vee I$ 2 3
1,2	5	\perp	$\Rightarrow E$ 1, 4
1	6	Hence $\neg A$	$\Rightarrow I$ 2, 5
1	7	$\neg A \vee B$	$\vee I$ 6

Forth exemple (example 3.1.12)

Prove $\neg A \vee B$ in the environment $A \Rightarrow B$.

environment			
reference		formula	
<i>i</i>		$A \Rightarrow B$	
context	number	proof	justification
1	1	Suppose $\neg(\neg A \vee B)$	
1,2	2	Suppose A	
1,2	3	B	$\Rightarrow E\ i, 2$
1,2	4	$\neg A \vee B$	$\vee I\ 2\ 3$
1,2	5	\perp	$\Rightarrow E\ 1, 4$
1	6	Hence $\neg A$	$\Rightarrow I\ 2, 5$
1	7	$\neg A \vee B$	$\vee I\ 1\ 6$
1	8	\perp	$\Rightarrow E\ 1, 7$

Forth exemple (example 3.1.12)

Prove $\neg A \vee B$ in the environment $A \Rightarrow B$.

environment			
reference		formula	
<i>i</i>		$A \Rightarrow B$	
context	number	proof	justification
1	1	Suppose $\neg(\neg A \vee B)$	
1,2	2	Suppose A	
1,2	3	B	$\Rightarrow E$ i, 2
1,2	4	$\neg A \vee B$	$\vee I$ 2 3
1,2	5	\perp	$\Rightarrow E$ 1, 4
1	6	Hence $\neg A$	$\Rightarrow I$ 2, 5
1	7	$\neg A \vee B$	$\vee I$ 6
1	8	\perp	$\Rightarrow E$ 1, 7
	9	Hence $\neg\neg(\neg A \vee B)$	$\Rightarrow I$ 1, 9

Forth exemple (example 3.1.12)

Prove $\neg A \vee B$ in the environment $A \Rightarrow B$.

environment			
reference		formula	
i		$A \Rightarrow B$	
context	number	proof	justification
1	1	Suppose $\neg(\neg A \vee B)$	
1,2	2	Suppose A	
1,2	3	B	$\Rightarrow E$ 1, 2
1,2	4	$\neg A \vee B$	$\vee I$ 2 3
1,2	5	\perp	$\Rightarrow E$ 1, 4
1	6	Hence $\neg A$	$\Rightarrow I$ 2, 5
1	7	$\neg A \vee B$	$\vee I$ 6
1	8	\perp	$\Rightarrow E$ 1, 7
	9	Hence $\neg\neg(\neg A \vee B)$	$\Rightarrow I$ 1, 9
	10	$\neg A \vee B$	RAA 9

Tree (example 3.1.12)

Give the **tree** representation of the previous proof :

Tree (example 3.1.12)

Give the **tree** representation of the previous proof :

$$\begin{array}{c}
 \frac{(i)A \Rightarrow B \quad (2)\cancel{A}}{(3)B} \Rightarrow E \\
 \frac{(1)\cancel{\neg(\neg A \vee B)} \quad \frac{(4)\neg A \vee B}{\vee 2}}{\Rightarrow E} \\
 \frac{(5)\perp}{(6)\neg A} \Rightarrow I[2] \\
 \frac{(1)\cancel{\neg(\neg A \vee B)} \quad \frac{(7)\neg A \vee B}{\vee 1}}{\Rightarrow E} \\
 \frac{(8)\perp}{(9)\neg\neg(\neg A \vee B)} \Rightarrow I[1] \\
 \frac{(9)\neg\neg(\neg A \vee B)}{(10)\neg A \vee B} \text{RAA}
 \end{array}$$

Tree (example 3.1.12)

Give the **tree** representation of the previous proof :

$$\begin{array}{c}
 \frac{(i)A \Rightarrow B \quad (2)\cancel{A}}{(3)B} \Rightarrow E \\
 \frac{(1)\cancel{\neg(\neg A \vee B)} \quad \frac{(4)\neg A \vee B}{\vee 2}}{\Rightarrow E} \\
 \frac{(5)\perp}{(6)\neg A} \Rightarrow I[2] \\
 \frac{(1)\cancel{\neg(\neg A \vee B)} \quad \frac{(7)\neg A \vee B}{\vee 1}}{\Rightarrow E} \\
 \frac{(8)\perp}{(9)\neg\neg(\neg A \vee B)} \Rightarrow I[1] \\
 \frac{(9)\neg\neg(\neg A \vee B)}{(10)\neg A \vee B} \text{RAA}
 \end{array}$$

The environment consists of formulae occurring at non-removed leaves.

Conclusion

Thank you for your attention.

Questions ?